


Things you should know when you leave Discussion today for one-electron atoms:

- $E_n = -R_y \frac{Z^2}{n^2} = -2.179 \cdot 10^{-18} \text{J} \frac{Z^2}{n^2} = -13.6 \text{eV} \frac{Z^2}{n^2}$
- $\Delta E_{\text{matter}} = E_n - E_m$; **Ionization Energy (IE)** = $E_\infty - E_{n(\text{initial})}$
- $\Delta E_{\text{light}} = h\nu_{\text{light}} = \text{IE} + \text{KE}$


1. Consider the following energy levels of the hydrogen atom as shown below in the diagram:

- Sketch electron clouds corresponding to energy levels up to $n=3$. (next to the lines on the right \rightarrow)
- When electron cloud of the H atom at the lowest energy interacts with light of the lowest resonate frequency, sketch the resulting electron cloud(s).
- When electron cloud of the H atom at the lowest energy interacts with light of the 2nd lowest resonate frequency, sketch the resulting electron cloud(s).
- When electron cloud of the He atom at the lowest energy interacts with light of the lowest resonate frequency, sketch the resulting electron cloud(s).
- What are the two energy levels involved in the ionization energy of an electron from H in the ground state?
- What is the ground state H atom ionization energy in J, eV and kJ/mol?
($\text{H}(g) \rightarrow \text{H}^+(g) + e^-$)



- What is the expression for the ionization energy of an electron from the $n=3$ state of an atom with atomic number Z ?

2. Hypothetical atom, X, has the following **ground state absorption** spectrum (displayed below), and the ground state ionization energy of the atom is 15eV (IE). Each of the **absorption** lines corresponds to a natural frequency of the electron cloud resulting from the mixing of electron wave with 1 loop with electron wave with more than one loop (i.e. the atom always starts in its **ground state** in each **absorption**). (In answering the following questions, assume that the zero of energy is the ionized atom)
- $$X \rightarrow X^+ + e^-.$$




- (a) What is the energy of the ground state (of the electron cloud corresponding to one loop) of X? (Careful of a sign!)
- (b) Construct the energy level diagram for X. (Hint: How many energy levels are necessary to account for the **absorption** spectrum?)
- (c) Using the energy level diagram you just did draw: How many lines will be present in the electrically-excited emission spectrum of this atom?
- (d) What is the energy of the *third excited* state of X? How many loops does that state have?
- (e) What is the energy of the *highest energy state* of X that is necessary to account for the **absorption** spectrum?
- (f) Is it possible for an electron in the ground state of atom X to absorb light of energy 3 eV or will it be transparent to it?

- (g) Can atom X emit light of energy 3 eV?
- (h) Is it possible for X to absorb light of energy 16eV or will it be transparent to it? Why?
- (i) Can atom X emit light of energy 16 eV?
- (j) What is the lowest frequency of the light emitted from the atom?
- (k) Draw the emission spectrum:

3. What is the expression for the energy in J for the ground state of an electron cloud in Li^{2+} ?

- a. Calculate wavelength (in nm) of the light corresponding to the Li^{2+} electron cloud resulting from mixing a 3 loop electron wave with a 1 loop electron wave? (Answer: 11nm)

4. What electron cloud energies account for the line corresponding to the wavelength 434 nm in the gas discharge spectrum for the Balmer series of H atoms? (In the Balmer series in each of the emission lines corresponds to a natural frequency of the electron cloud resulting from the mixing of an electron wave with more than two loop with the electron wave with 2 loop)



- A. Only the $n = 3$ cloud energy
 - B. Only the $n = 4$ cloud energy
 - C. Only the $n = 5$ cloud energy
 - D. The $n = 2$ and $n = 4$ cloud energies
 - E. The $n = 2$ and $n = 5$ cloud energies
 - F. None of these.
5. Photons of energy $13.6\text{eV} = R_y$ are able to ionize H in its $n = 1$ energy level. Are photons of this energy able to ionize He^+ in its $n = 2$ energy level?

6. The light with wavelength of 365 nm will ionize H atom in the $n = 2$ energy level. What effect will light with wavelength = 657 nm have? (Choose all that apply)
- Atom will be transparent to the light of this wavelength.
 - Ionization will take place
 - Ionization will not take place
 - The ionized electron will use excess energy for kinetic energy
7. The light with wavelength of 365nm will ionize H atom in the $n = 2$ energy level. What effect will light with wavelength = 265nm have? (Choose all that apply)
- Atom will be transparent to the light of this wavelength.
 - Ionization will take place
 - Ionization will not take place
 - The ionized electron will use excess energy for kinetic energy
8. The photoelectric effect threshold frequency of a metal is $\nu_0 = 1 \cdot 10^{15}$ Hz. Gamma radiation of frequency $1 \cdot 10^{17}$ Hz ejects electrons from the metal. Which of the following occurs when the intensity of the gamma radiation is reduced by 50 %?
- The velocity of the ejected electrons will be reduced by a factor of two.
 - The kinetic energy of the ejected electrons will be reduced by a factor of two.
 - The kinetic energy of the ejected electrons will be reduced by a factor of four.
 - Kinetic energy and the velocity of the ejected electrons will stay the same.
 - Number of ejected electrons will increase.
 - Number of ejected electrons will decrease.
9. Assume light is able to eject electrons from a metal. What do you expect as the wavelength of the light is increased?
- If the light wavelength reaches the lowest, electrons will no longer be ejected.
 - Electrons will still be ejected but they will move faster and faster.
 - Electrons will still be ejected but they will move slower and slower.
 - More and more electrons will be ejected but they will have the same kinetic energy.
 - More information needed.
10. An electron is ionized from the ground state of an atom, with $E_1 = -1.3 \cdot 10^{-19}$ J, by light with frequency $1.0 \cdot 10^{15}$ Hz. What is the kinetic energy (in eV) of the ejected electron?

11. The work function (ionization energy), of chromium metal is 7.2×10^{-19} J. What is the maximum kinetic energy of an electron, if it is ejected from chromium metal by light of wavelength 250.nm?
- What will happen to the speed and quantity of the ejected electrons if the wavelength of the light will increase?
 - What will happen to the speed and quantity of the ejected electrons if the wavelength of the light will decrease?
 - What will happen to the speed and quantity of electrons if the intensity of the light will increase?
12. An atom has only three energy levels, -2.5, -4.0, and -4.5 eV. Draw the **absorption** spectrum for a gas of these atoms excited by an electric discharge.
(Hint: you need to draw energy level diagrams to answer this question assume the most negative energy is an energy of the ground state)
- Draw the **emission** spectrum. (Hint: does it always just goes to the ground state?)
13. An atom has only three energy levels, -2.6, -4.1, and -5 eV. Draw the **absorption** spectrum for a gas of these atoms excited by an electric discharge.
- List all the lines in the **emission** spectrum. Do not assume that it always goes to the ground state.
14. One atom emits light of energy 1.27 eV. A second atom has only three energy levels: -0.9 eV, -1.27 eV, and -1.9 eV. Assuming ionization is not possible can the second atom absorb the light emitted by the first atom.


Useful information:

- Atomic wave function family album
 - a. Principle quantum number $n=1,2,3,\dots$ specifies energy level
 $n = j + l$
 - b. Number of nodal planes (l).
 - c. Angular momentum quantum number ($l=0, 1, 2,\dots n-1$) defines the shape of the orbital.
 - d. $m_l = -l, \dots, 0, \dots, l$
 - e. Number of loops j ($j=n-l$)
 - f. S, P and D and F orbitals.
 - g. Size of the orbital is proportional to n^2
- Electron configuration
 - a. Orbital, Shell, Subshell
 - b. Shielding
 - c. Pauli Exclusion Principle. Auf-bau. Hund's rule
- Periodic Trends (**I.E.**, E.A., Radius, Ionic Radius, Electronegativity)
- For many electron atoms:

$$E_n = -13.6(\text{eV}) \frac{Z_{\text{eff}}^2}{n^2} = -2.18 \cdot 10^{-18} (\text{J}) \frac{Z_{\text{eff}}^2}{n^2} = -1312 \left(\frac{\text{kJ}}{\text{mol}} \right) \frac{Z_{\text{eff}}^2}{n^2}$$


Useful Information

- Below $3d_{xy}$, $3d_{x^2-y^2}$ and $3d_{yz}$ orbital. Be sure to correctly orient your orbitals in the x-y-z space.




Orbital	l	$n=j+l$	$j=n-l$	Shape	Number of orbitals in a subshells(m_l) $2l+1$	# e^- that can fit
s	0	1 2 3 etc	1 2 3 $j=n$	Sphere	1 orientation in space	2
p	1	2 3 4 etc	1 2 3 $j=n-1$	Dumbbell	3	6
d	2	3 4 etc	1 2 $j=n-2$	Cloverleaf	5	10
f	3	4 5 etc	1 2 $j=n-3$		7	14

1. For the two hydrogen electron clouds below identify the quantum number “ n ”, the quantum number “ l ”, the number of radial loops “ j ”, and the specific name of the orbital (you must indicate orientation, e.g. $3d_{xy}$).



- a. For the two hydrogen electron clouds above, which has the largest ionization energy? (Note: in H atom, all orbitals of same n have the same energy.)
2. What are the possible angular momentum quantum numbers for an orbital in the $n=4$ shell?
- How many **degenerate** orbitals are in the $n=3$ shell? What are they? (Note: in H atom, all orbitals of same n have the same energy.)

3. The figure shows three H atom electron clouds.
- Which of the following H atom electron clouds has the largest ionization energy?
 - Write down the **numerical expression** that when evaluated gives the value **in eV** of **smallest** ionization energy of these three clouds. (Answer: 0.85)



Additional Practice problems:

- The ionization wavelength of H atom in the $n = 2$ energy level is 365 nm. Will light of this wavelength ionize He^+ in the $n = 1$ level?
- An electron is ionized from the ground state of an atom, with $E_1 = -30$ eV, by light for which a photon of light energy is 36 eV. What is the kinetic energy (in eV) of the ejected electron? (Answer: 6eV)
- What is kinetic energy of the electron ionized from Li^{2+} in its $n = 6$ level by light of wavelength 310 nm? (Answer: 0.94×10^{-19})
- In terms of the variables (like: R_y , h , and c) what is the smallest possible wavelength of light that will be emitted by a He^+ atom starting with the energy corresponding to 6 loops?
- What is the largest possible wavelength of light that will be emitted by the Li^{2+} atom with the energy level corresponding to 6 loops? (Answer: 800nm)

In preparation for lectures and next week discussion:

Please bring Page 9 and 10 below to the lectures until completed

Atom	Z	Electron configuration	IE_1 $\frac{kJ}{mol}$	Z_{eff}	Trends in IE_1 is Explained by: a. Z increases b. Electron –electron repulsion c. New shell d. l increases or Z_{eff} decreases (shielding)	Ions (Ions Electron configuration)
He	2	$1s^2$	2373			He^+ $1s^1$
Li	3	$1s^2 2s^1$	520			Li^+ $1s^2$
Be	4	$1s^2 2s^2$	899			Be^+ $1s^2 2s^1$
B	5		801			B^+
C	6	$1s^2 2s^2 2p_x^1 2p_y^1$	1086			C^+ $1s^2 2s^2 2p_x^1$
N	7		1400			N^+ $1s^2 2s^2 2p_x^1 2p_y^1$
O	8		1314			O^+ $1s^2 2s^2 2p_x^1 2p_y^1 2p_z^1$
F	9	$1s^2 2s^2 2p_x^2 2p_y^2 2p_z^1$	1680			F^+
Ne	10		2080			Ne^+ $1s^2 2s^2 2p_x^2 2p_y^2 2p_z^1$
Na	11		496			Na^+ $1s^2 2s^2 2p_x^2 2p_y^2 2p_z^2$
Mg	12	$1s^2 2s^2 2p_x^2 2p_y^2 2p_z^2 3s^2$ = $[Ne]3s^2$	738			Mg^+ $[Ne]3s^1$
Al	13	$[Ne]3s^2 3p_x^1$	578			Al^+ $[Ne]3s^2$

Atom	Z	Ions (Ions Electron configuration)	IE ₂ $\frac{kJ}{mol}$	Z _{eff} For the Ions	Trends in IE ₂ is Explained by: a. Z increases b. Electron –electron repulsion c. New shell d. l increases or Z _{eff} decreases (shielding)
He	2	He ⁺ 1s ¹	5248		
Li	3	Li ⁺ 1s ²	7300		
Be	4	Be ⁺ 1s ² 2s ¹	1757		
B	5	B ⁺	2430		
C	6	C ⁺ 1s ² 2s ² 2p _x ¹	2350		
N	7	N ⁺ 1s ² 2s ² 2p _x ¹ 2p _y ¹	2860		
O	8	O ⁺ 1s ² 2s ² 2p _x ¹ 2p _y ¹ 2p _z ¹	3390		
F	9	F ⁺	3370		
Ne	10	Ne ⁺ 1s ² 2s ² 2p _x ² 2p _y ² 2p _z ¹	3950		
Na	11	Na ⁺ 1s ² 2s ² 2p _x ² 2p _y ² 2p _z ²	4560		
Mg	12	Mg ⁺ [Ne]3s ¹	1450		
Al	13	Al ⁺ [Ne]3s ²	1820		

6. Draw the electron configuration diagram for C, but **fill** the diagram as if the C is in the ground state. Draw a dotted line around a shell. Circle a subshell. Draw a triangle around an orbital. Is C atom paramagnetic or diamagnetic?

a. Calculate Z_{eff} of C, if $IE_1 = 1086 \text{ kJ/mol}$. (Hint: $IE = E_{\infty} - E_{\text{initial}} = 2.18 \cdot 10^{-18} (\text{J}) \frac{Z_{\text{eff}}^2}{n^2}$)

Exam 3 answers:

1. $-\Delta H(\text{BA}) + \Delta H(\text{BE}) + \Delta H(\text{EC})$

2.

a. For water, B is $<$ than D.

b. For water, slope of A is $>$ than slope of C.

c. For water, slope of C is $<$ than slope of E.

d. B for water is $>$ B for ammonia (NH_3).

e. D for methane (CH_4) is $<$ D for methanol (CH_3OH).

3.

a. -188 kJ/mol

b. -1636 kJ/mol

c. -42 kJ

d. $50. \text{ kJ.mol}$


e. 6 kJ/mol

4.

a. $5.11 \cdot 10^{-19} \text{ J}$

b. 0.61

5. $19.6 \text{ }^\circ\text{C}$



7.

a. -318 kJ/mol

b. $q_p < 0$ $w > 0$ $\Delta U < 0$

$T_{\text{final}} (\text{constant pressure}) > T_{\text{final}} (\text{constant volume})$

8. $\text{C}_2 < \text{C}_3 < \text{C}_1$

9. F, F, F, T, F

10. $3.76 \cdot 10^{13} \text{ Hz}$

11. 74.5%