

Lecture 27 CH102 A1 (MWF 9 am) Spring 2017 Copyright © 2016 Dan Dill dan@bu.edu

[TP] A redox reaction has $K=10$. If there are **only reactants** present, $-\log(Q/K)$ is ...

17% 1. $+\infty$
 17% 2. 2
 17% 3. 1
 17% 4. 0
 17% 5. -1
 17% 6. $-\infty$

BOSTON UNIVERSITY Response Counter 10 1

Lecture 27 CH102 A1 (MWF 9:05 am)
 Monday, April 3, 2017

- Complete: Cell voltage versus spontaneity.
- The Nernst equation.

Next lecture: Complete ch16: Exploring the Nernst equation.
 Concentration cells: Mixing \rightarrow electric current

BOSTON UNIVERSITY

Lecture 27 CH102 A1 (MWF 9 am) Spring 2017 Copyright © 2016 Dan Dill dan@bu.edu

[TP] A redox reaction has $K=10$. If there are **only reactants** present, $-\log(Q/K)$ is ...

17% 1. $+\infty$
 17% 2. 2
 17% 3. 1
 17% 4. 0
 17% 5. -1
 17% 6. $-\infty$

BOSTON UNIVERSITY Response Counter 10 8

Lecture 27 CH102 A1 (MWF 9 am) Spring 2017 Copyright © 2016 Dan Dill dan@bu.edu

[Quiz] For the redox process
 $\text{Cu}^{2+}(aq) + \text{Zn}(s) \rightarrow \text{Cu}(s) + \text{Zn}^{2+}(aq)$
 when $[\text{Cu}^{2+}] = 0.001$ and $[\text{Zn}^{2+}] = 0$, the voltage is ...

33% 1. < 0
 33% 2. 0
 33% 3. > 0

BOSTON UNIVERSITY Response Counter 10 9

Lecture 27 CH102 A1 (MWF 9 am) Spring 2017 Copyright © 2016 Dan Dill dan@bu.edu

Voltage E versus Q/K

Make a table of Q , Q/K and $-\log(Q/K)$ for a reaction with $K=10$, for values $Q=0, 0.1, 1, 10, 100$, and ∞ .

Q	Q/K	$-\log(Q/K)$
0		
0.1		
1		
10		
100		
∞		

BOSTON UNIVERSITY

10

Lecture 27 CH102 A1 (MWF 9 am) Spring 2017 Copyright © 2016 Dan Dill dan@bu.edu

Voltage E versus Q/K

Make a table of Q , Q/K and $-\log(Q/K)$ for a reaction with $K=10$, for values $Q=0, 0.1, 1, 10, 100$, and ∞ .

Q	Q/K	$-\log(Q/K)$
0	0	$+\infty$
0.1	0.01	2
1	0.1	1
10	1	0
100	10	-1
∞	∞	$-\infty$

BOSTON UNIVERSITY

11

Lecture 27 CH102 A1 (MWF 9 am) Spring 2017 Copyright © 2016 Dan Dill dan@bu.edu

Voltage E versus Q/K

Using your table, sketch $-\log(Q/K)$ versus Q/K , with $K=10$, for values $Q=0, 0.1, 1, 10, 100$, and ∞ .

Q	Q/K	$-\log(Q/K)$
0	0	$+\infty$
0.1	0.01	2
1	0.1	1
10	1	0
100	10	-1
∞	∞	$-\infty$

BOSTON UNIVERSITY

13

Lecture 27 CH102 A1 (MWF 9 am) Spring 2017 Copyright © 2016 Dan Dill dan@bu.edu





Voltage E versus Q/K

Using your table, plot $-\log(Q/K)$ versus Q/K , with $K=10$, for values $Q=0, 0.1, 1, 10, 100$, and ∞ .

Q	Q/K	$-\log(Q/K)$
0	0	$+\infty$
0.1	0.01	2
1	0.1	1
10	1	0
100	10	-1
∞	∞	$-\infty$

BOSTON UNIVERSITY

14



Lecture 27 CH102 A1 (MWF 9 am) Spring 2017

Copyright © 2016 Dan Dill dan@bu.edu

$$E = -(0.06/n_e) \text{ V } \log(Q/K)$$

Calculate the voltage at 25 °C for $n_e = 1$ when $Q = (10) \times K$

$$E = -0.06 \text{ V}$$



19

Lecture 27 CH102 A1 (MWF 9 am) Spring 2017

Copyright © 2016 Dan Dill dan@bu.edu

$$E = -(0.06/n_e) \text{ V } \log(Q/K)$$

At 25 °C for $n_e = 1$, ...
each **order of magnitude** change in Q/K ...
changes voltage by **0.06 V**.



20

Lecture 27 CH102 A1 (MWF 9 am) Spring 2017

Copyright © 2016 Dan Dill dan@bu.edu

$$E = -(0.06/n_e) \text{ V } \log(Q/K)$$

Write an expression for E when $Q = 1$.



21

Lecture 27 CH102 A1 (MWF 9 am) Spring 2017

Copyright © 2016 Dan Dill dan@bu.edu

$$E = -(0.06/n_e) \text{ V } \log(Q/K)$$

The value of E when $Q = 1$ is called the **standard voltage** and at 25 °C is written as

$$E(Q = 1) = E^\circ = +(0.06/n_e) \text{ V } \log(K)$$



22

Lecture 27 CH102 A1 (MWF 9 am) Spring 2017

Copyright © 2016 Dan Dill dan@bu.edu

[TP] The value of E when $Q = 1$ at 25 °C is

$$E(Q = 1) = E^\circ + (0.06/n_e) V \log(K)$$

For $n_e = 1$, if K is different by a factor of ten (say, 17 instead of 1.7), the magnitude of standard voltage will change by ...

- 20% 1. 10 V
- 20% 2. 1 V
- 20% 3. 0.1 V
- 20% 4. 0.06 V
- 20% 5. Some other amount



23

Lecture 27 CH102 A1 (MWF 9 am) Spring 2017

Copyright © 2016 Dan Dill dan@bu.edu

[TP] The value of E when $Q = 1$ at 25 °C is

$$E(Q = 1) = E^\circ + (0.06/n_e) V \log(K)$$

For $n_e = 3$, if K is different by a factor of ten (say, 17 instead of 1.7), the magnitude of standard voltage will change by ...

- 25% 1. 0.18 V
- 25% 2. 0.06 V
- 25% 3. 0.02 V
- 25% 4. Some other amount



24

Lecture 27 CH102 A1 (MWF 9 am) Spring 2017

Copyright © 2016 Dan Dill dan@bu.edu

[Quiz] The value of E when $Q = 1$ at 25 °C is

$$E(Q = 1) = E^\circ + (0.06/n_e) V \log(K)$$

A typical physiological value of E° is 0.18 V.

For $n_e = 1$ this corresponds to the value of K equal to ...

- 17% 1. 0.1
- 17% 2. 1
- 17% 3. 10
- 17% 4. 100
- 17% 5. 1000
- 17% 6. Some other value



25

Lecture 27 CH102 A1 (MWF 9 am) Spring 2017

Copyright © 2016 Dan Dill dan@bu.edu

$$E = -(0.06/n_e) V \log(Q/K)$$

The value of E when $Q = 1$ at 25 °C is

$$E(Q = 1) = E^\circ + (0.06/n_e) V \log(K)$$

Calculate K corresponding to $E^\circ = 1.8$ V for $n_e = 1$.

$K = 10^{30}$. Very large!



26

Lecture 27 CH102 A1 (MWF 9 am) Spring 2017

Copyright © 2016 Dan Dill dan@bu.edu

$$E = -(0.06/n_e) \text{ V } \log(Q/K)$$

The value of E when $Q = 1$ at 25 °C is

$$E(Q = 1) = E^\circ = +(0.06/n_e) \text{ V } \log(K)$$

Express the cell voltage for **any value of Q** in terms of E° , that is, in terms of the cell voltage when $Q = 1$.



27

Lecture 27 CH102 A1 (MWF 9 am) Spring 2017

Copyright © 2016 Dan Dill dan@bu.edu

$$E = -(0.06/n_e) \text{ V } \log(Q/K)$$

The value of E when $Q = 1$ at 25 °C is

$$E(Q = 1) = E^\circ = +(0.06/n_e) \text{ V } \log(K)$$

The cell voltage at 25 °C for **any value of Q** in terms of the cell voltage when $Q = 1$ is

$$E(\text{any } Q) = E^\circ - (0.06/n_e) \text{ V } \log(Q)$$

This is called the **Nernst equation**



28