

**Things you should know when you finish the Discussion hand out:**

1. Gas Laws  $P\Delta V_{\text{gas}} = n_{\text{gas}}RT$

$$T = T [^{\circ}\text{C}] + 273.15 [\text{K}]$$

$$\begin{aligned} R &= 8.314 [\text{J}/(\text{mol}\cdot\text{K})] = 8.314 [(\text{L}\cdot\text{kPa})/(\text{mol}\cdot\text{K})] = 0.08206 [(\text{L}\cdot\text{atm})/(\text{mol}\cdot\text{K})] \\ &= 0.08314 [(\text{L}\cdot\text{bar})/(\text{mol}\cdot\text{K})] \end{aligned}$$

2. 
$$\bar{E}_k = \frac{M\bar{u}^2}{2} = \frac{3}{2}RT;$$

3. 
$$\bar{u}^2 = \frac{3RT}{M}$$

4. 
$$\text{rate} = \frac{1}{\text{time}} = \frac{\text{amount}}{\text{time}}; \text{ rate} \sim \text{velocity}$$

1. Sketch the MO diagram for  $\text{C}_4\text{H}_2$ , ( $\text{C}_A\text{H}\equiv\text{C}_B-\text{C}_C\equiv\text{C}_D\text{H}$ ).a. Determine and draw  $\sigma$  framework:b. How many pairs of electrons are in the  $\sigma$  frameworkc. How many pairs of electrons are in the  $\pi$  framework?d. Draw correlation diagram for  $\pi$  framework:

- e. How many pairs of electrons are shared between the middle two carbons in  $C_4H_2$ ,
- f. How many pairs of electrons are shared between the first two carbons in  $C_4H_2$ ,
2. A bicycle tire is filled with air to a pressure of 4 atm, at a temperature of  $27^\circ C$ . Riding the bike on asphalt on a hot day increased the temperature of the tire to ( $78^\circ C$ ). The volume of the tire has increased by 4.0%.  
Draw a picture of the tire before and after the ride, showing variables associated with the tire in each situation. (Hint: Find mathematical relationship between initial and final volume)

What is the new pressure in the bike tire?

Do you need a new volume of the tire after the ride when solving for the new pressure?

Can you determine the volume of the tire after the ride?

If not, what other information do you need?

3. Air bags are activated when a severe impact causes a steel ball to compress a spring and electrically ignite a detonator cap. This action causes sodium azide ( $NaN_3$ ) to decompose explosively according to the equation below:



What mass of sodium azide must be reacted to inflate an air bag to 100. L at  $27^\circ C$  and 1 atm?  
(Answer: 200g)

4. A 20 L stainless steel container was filled with 2 atm of  $\text{H}_2$  gas and 3 atm of  $\text{O}_2$  gas. A spark ignited the mixture, producing water.
- What is the pressure in the tank if the reaction is run at  $125^\circ\text{C}$ ?
  - What is the pressure in the tank if the reaction is run at  $25^\circ\text{C}$ ?
- (Hint: Think about the state of water at those temperatures)(Answer: 2atm, 4atm)
- c. Challenge question (**to do at home**) How will liquid water affect the volume of the container and the pressure?
5. At STP, 1 L of  $\text{Br}_2$  reacts with 3 L of  $\text{F}_2$  to completion and produces 2 L of a product. What is the formula of the product? (All substances are gases. Hint: 1 mol of all gasses occupy the same volume)
6. A rigid 4.0 L container contains 0.20 mol of  $\text{H}_2$  and 0.20 mol of  $\text{O}_2$  at 300. K. Circle your answer to each of the following questions.
- On average, molecules of which gas hit the container walls more times per second?  

$\text{H}_2$	$\text{O}_2$	the same	further information required
--------------	--------------	----------	------------------------------
  - On average, molecules of which gas hit the container walls with greater momentum?  

$\text{H}_2$	$\text{O}_2$	the same	further information required
--------------	--------------	----------	------------------------------
  - On average, molecules of which gas hit the container walls with greater force?  

$\text{H}_2$	$\text{O}_2$	the same	further information required
--------------	--------------	----------	------------------------------
  - On average, molecules of which gas contribute more to the total pressure?  

$\text{H}_2$	$\text{O}_2$	the same	further information required
--------------	--------------	----------	------------------------------
  - What is the total pressure exerted on the walls of the container, in atm?(Answer:2.5 atm)

7. Which of the following combinations of conditions will a gas behave most ideally? Explain your choice.
- Low P and low T
  - Low P and high T
  - High P and high T
  - High P and low T
8. Consider two identical cylinders, one containing 1.0 mole of He and the other containing 1.0 mole of N<sub>2</sub>. The cylinder of nitrogen gas has a constant temperature of 77°C.
- Will He atoms need to be hotter or colder to have the same rms speed as N<sub>2</sub> gas? Why? (*Hint: Think about this conceptually. Do not try to plug numbers in*)
  - Calculate the temperature in K of the He such that it will have the same rms speed as the N<sub>2</sub> gas at 77°C. (**Hint: You do not need to calculate the value of rms speed to solve this problem.**) ( Answer: 50K)
  - If both gases act ideally, circle which of the following must be true about the average kinetic energy, and gas pressure if both gases are at the same T? (Hint: think what is the pressure and force depends on?)
 

$\bar{E}_{He} > \bar{E}_{N_2}$	$\bar{E}_{He} < \bar{E}_{N_2}$	$\bar{E}_{He} = \bar{E}_{N_2}$
$P_{He} > P_{N_2}$	$P_{He} < P_{N_2}$	$P_{He} = P_{N_2}$
  - If identical holes are opened in each cylinder, what will be true about the time for the gases to effuse out their cylinders if both gases are at the same T?(Hint: faster means less time spend)
 

time <sub>(He)</sub> > time <sub>(N<sub>2</sub>)</sub>	time <sub>(He)</sub> < time <sub>(N<sub>2</sub>)</sub>	time <sub>(He)</sub> = time <sub>(N<sub>2</sub>)</sub>	More Information Needed
--	--	--	-------------------------
9. At 0 °C and 1 bar = 100 kPa, the average speed of oxygen molecules is **460 m/s**. When the pressure is **lowered by half** at 0 °C, the average speed will be:
10. A sealed balloon is filled with 1 L of helium at 27°C and 1 atm. The balloon rises to a point in the atmosphere where the pressure is 0.3 atm and the temperature is -33 °C. What is the density inside the balloon after the balloon rises ( $M = \frac{mRT}{VP} = \frac{\rho RT}{P}$ )? (At home: What is the density inside the balloon before it rises?) (Hint: Will the density be the same or different at the beginning and at the end?) (Answer:0.16, 0.0609)

First predict will balloon expand or not?

What is the volume of the balloon at the end? ( Answer:3L)

11. Methanol,  $\text{CH}_3\text{OH}$ , can be produced by the following reaction:  $\text{CO}(\text{g}) + 2\text{H}_2(\text{g}) \rightarrow \text{CH}_3\text{OH}(\text{g})$ . Hydrogen at SATP flows into a reactor at a rate of 12.5 L/min. Carbon monoxide at SATP flows into the reactor at a rate of 25 L/min. If 6.4 g of methanol is produced per minute, what is the percent yield of the reaction per minute? (Hint: solve considering that everything happened in 1 minute.)(SATP  $T=25^\circ\text{C}$  ,  $P = 1\text{bar} = 100\text{kPa}$ ,  $V$  (of 1 mole of the gas) =  $\frac{1 \cdot R \cdot T}{P} = 24.789\text{L} = 25\text{L}$ ) ( Answer: 80%)

12. At  $20^\circ\text{C}$  the average speed of oxygen molecules is 500 m/s. What will be the average speed of xenon atoms (Answer: 250) :


13. The balanced equation for combustion of ethane at 400. K is



When 4.0 mol of ethane and 20. mol of oxygen are combined in a rigid container at 400. K, the total pressure is 4.8 atm. A spark ignites the mixture, resulting in complete combustion. Calculate the final pressure at 400. K.(Answer: 5.2atm)

14. At a certain temperature,  $T_X$ , gas X (molar mass  $M_X$ ) has an rms speed of 805 m/s. Derive an expression in terms of  $T_X$  for the temperature,  $T_Y$ , at which substance Y (molar mass  $M_Y$ ) has the same rms speed. Write your expression in the box.


15. The graph shows the distribution of speeds of ozone,  $\text{O}_3$ , at 1200 K. Sketch on the graph the distribution of speeds of methane,  $\text{CH}_4$ , at 400 K.



16. A mixture of 1.0 g of hydrogen gas ( $\text{H}_2(\text{g})$ ) and 1.0 g of helium gas ( $\text{He}(\text{g})$ ) exerts a pressure of 0.50 atm. What is the partial pressure of each gas in the mixture? (Answers: 0.33, 0.16, 66%, 33%)

$$(n_{\text{total}} = n_1 + n_2 + n_3; p_{\text{total}} = p_1 + p_2 + p_3 = (n_1 + n_2 + n_3) \frac{RT}{V}; \quad \chi_1 = \frac{n_1}{n_{\text{total}}}; \quad p_1 = \chi_1 \cdot p_{\text{total}})$$

17. Hydrogen chloride gas produced in a 100. L container according to the following reaction:



Initially 5.00 moles of  $\text{H}_2$  and 3.00 moles of  $\text{Cl}_2$  react. After the reaction is complete, the pressure in the reactor is 3.00 atm. Assuming that the process went to completion, what is the partial pressure (in atm) of all the gases in the container? (Answers: 0.750, 2.25)

What is the percent of each gas in the mixture?

18. The Haber process is used to produce gaseous ammonia from gaseous  $\text{N}_2$  and  $\text{H}_2$ . A 10 L reactor is charged with 28 g of  $\text{N}_2$  and 3 g of  $\text{H}_2$ . After the reaction is complete, the pressure in the reactor is 3 atm. Assuming that the process went to completion, what is the partial pressure (in atm) of ammonia in the reactor? (Answers: 2 atm)

19. A mixture of gases contains  $n_A$  moles of  $\text{CH}_4$ ,  $n_B$  moles of  $\text{C}_2\text{H}_6$ , and  $n_C$  moles of  $\text{C}_3\text{H}_8$ . The total pressure is  $P_T$ . Derive an expression, in terms of only the variables listed in this problem, for the partial pressure of  $\text{CH}_4$ ,  $P_A$ .

20. Calcium is dissolved in the ocean and is used by marine organisms to form  $\text{CaCO}_3(\text{s})$  in skeletons and shells. When organisms die, their remains fall to the bottom. The amount of calcium carbonate that can dissolve in seawater depends on the pressure. This prevents the world's calcium from being tied up as insoluble  $\text{CaCO}_3(\text{s})$  at the bottom of the sea. At pressures of 414 atm, the shells slowly re-dissolve. Estimate the depth (in meters) where water exerts 414 atm of pressure **to one significant figure**. The density of sea water is  $1030 \text{ kg} \cdot \text{m}^{-3}$  and remember force is mass times the acceleration of gravity ( $F = mg$ ) where  $g = 9.81 \text{ m} \cdot \text{s}^{-2}$ . (Hint:  $P = F/A$  and  $A = V/h$ ,  $1 \text{ atm} = 101 \text{ kPa}$ ,  $1 \text{ Pa} = 1 \text{ kg}/(\text{m} \cdot \text{s}^2)$ ) (Answer: 4000m)

End of the chapter 11 suggested question: 38, 39 43, 45, 46, 49, 50, 53, 54, 56.