

Things you should know when you finish the Discussion hand out:

$$1. \bar{E}_k = \frac{m\bar{u}^2}{2} = \frac{3}{2}RT; \quad \bar{u}^2 = \frac{3RT}{M}$$

$$2. \text{Rate} = \frac{[\text{amount}]}{\text{time}}$$

3. Page 438 in the text book "Summary of Intermolecular Forces" Table 11.15.

$$4. \left(P_{\text{observed}} + a \left[\frac{n}{V} \right]^2 \right) (V_{\text{container}} - bn) = nRT$$

$$\text{Where } a \left[\frac{n}{V} \right]^2 \text{ - accounts for intermolecular forces; } a = \left[\frac{L^2 \text{bar}}{\text{mol}^2} \right] = \left[\frac{100 \text{kPa} \cdot L^2}{\text{mol}^2} \right]$$

$$bn \text{ - Accounts for non-zero molecule volume; } b = \left[\frac{L}{\text{mol}} \right]$$


5. Phase Diagram.

- a. Triple point
- b. Critical point

1. Two identical containers, one red and one yellow, are inflated with different gases at the same volume and pressure. Both containers have an identically sized hole that allows the gas to leak out. It takes four times as long for the yellow container to leak out compared to the red container. If the red container is twice as hot as the yellow container, what is the ratio of the molar masses of the gases ($M_{\text{yellow}} / M_{\text{red}}$). (Answer: 8)

2. A container is filled with He at 18 °C and 7.92 bar. It is found that the pressure of the container drops by 50% in 35 minutes, due to a small hole. If the container had instead been filled with Ne at 18 °C and 7.92 bar, what would be the pressure of the Ne after 35 minutes? Express your answer in bar. (Answer: 6.16)

3. Which of the following combinations of conditions will a gas behave most ideally? Explain your choice.
- Low P and low T
 - Low P and high T
 - High P and high T
 - High P and low
4. Solve the van der Waals real gas equation for the pressure.
- To investigate how the a coefficient affects the pressure, set the b coefficient to zero. Does a increase or decrease the pressure compared to what you would expect for an ideal gas?
 - To investigate how the b coefficient affects the pressure, set the a coefficient to zero. Does b increase or decrease the pressure compared to what you would expect for an ideal gas?
 - Below is a plot of the compressibility (PV/RT) as a function of changing pressure for 1 mole of a real gas (dashed line) and an ideal gas (solid line). Using what you learned in parts (a) and (b), at what pressures is the a coefficient dominant and at what pressures is the b coefficient dominant. (y axis is $(PV/RT)_{\text{real}} / (PV/RT)_{\text{ideal}}$)



5. Put the following molecules in order of increasing Van der Waals constant, b : Cl_2 , N_2 , C_2H_6 , C_3H_8 .

6. For each of the pair of molecules below, which has the highest value of Van der Waals constant, a ?
Identify the dominant intermolecular forces.

H₂O vs CO₂

Ne vs F₂,

C₆H₆ vs CH₃OH

7. Which of the following gases will behave least ideally under the same conditions? Explain your choice using the a and b Van der Waals coefficients.

CH₄ or SO₂

Cl₂ or N₂

8. Match the molecules below with their a and b van der Waals coefficients.

<u>Gas</u>	<u>a (L² atm mol⁻²) / b (L mol⁻²)</u>
Water (H ₂ O)	0.2476 / 0.02661
Argon (Ar)	5.536 / 0.03029
Hydrogen (H ₂)	1.363 / 0.03219
Benzene (C ₆ H ₆)	18.24 / 0.1154


9. O₂ and H₂O have similar values of van der Waals constant, b , but different van der Waals constant, a which one will have higher pressure at the same temperature and volume.
10. NO₂ and Trifluoromethane (CF₃H) have similar values of Van der Waals constant, a , but different van der Waals constant, b which one will have higher pressure at the same temperature and volume.
11. A 5.0mol sample of NH₃ gas is kept at 2.0 L container and 27°C, Calculate the pressure in bars of the gas assuming it does not behave ideally. ($a=4.0$ bar·L²/mol² and $b= 0.040$ L/mol)

What is the percent error compared to an ideal gas?

12. Consider a container filled with $^{15}\text{N}_2$ gas and placed in larger container filled only with $^{14}\text{N}_2$. If a small hole is made in the smaller container, initially as the gases effuse, will the smaller container gain or lose mass? *Assume the pressures inside the containers are the same.* Explain your answer. It may help to draw a picture.
13. Two identical containers, one red and one yellow, are filled with different noble gases at the same temperature and pressure. Both containers have an identically sized hole in them allowing the gases to leak out. If it takes the red container 10.0 seconds to empty and the yellow container 57.3 seconds to empty, what are the noble gases contained in each container? (Remember: rate is proportional to amount/time)
14. In 2.00 min, 29.7mL of helium effuses through a small hole. Under the same conditions (temperature and pressure), a 10.00 mL of a mixture of CO and CO₂ effuse through the same hole in the same amount of time. Calculate the percent composition by volume of the mixture. (Answer: 45.5% and 54.5%)
15. Two identical containers are filled with the same number of moles of gas at the same temperature. Container A contains a gas with a molar mass of 40. g/mol, and container B contains a gas with a molar mass of 160. g/mol. A hole of identical size is then made in each container. What is the ratio $\text{time}_A/\text{time}_B$ of the times necessary for the number of moles of gas to drop to half in each container? (Answer: 0.5)


16. Imagine a substance with the following points on the phase diagram: a triple point at 0.5 atm and -5°C ; a normal melting point at 20°C ; a normal boiling point at 150°C ; and a critical point at 5 atm and 1000°C . For this, complete the following:

- What does it mean to have a normal solid-liquid line?
- Roughly, sketch the phase diagram, using units of atmosphere and $T^{\circ}\text{C}$.



- Rank the states with respect to increasing density
- Describe what one would see at pressures and temperatures above 5 atm and 1000°C .
- Describe what will happen to the substance when it begins in a vacuum at -15°C and is slowly pressurized
- Describe the phase changes from -80°C to 500°C at 2 atm

17. For each of the following questions refer to the phase diagram for mysterious compound X.



- What is the critical temperature of compound X?
- If you were to have a bottle containing compound X in your closet, what phase would it most likely be in?
- At what temperature and pressure will all three phases coexist?
- If I have a bottle of compound X at a pressure of 45 atm and temperature of 100°C, what will happen if I raise the temperature to 400°C?
- Why can't compound X be boiled at a temperature of 200°C?
- If I wanted to, could I drink compound X?

The following will Not be part of exam 1 but will be on the quiz next week:

In Preparation for next week lecture and discussion sections:

- Using your own words define Lattice enthalpy ($\Delta_{\text{lattice}}H$) :
- Is the Lattice enthalpy ($\Delta_{\text{lattice}}H$) (choose one):

always positive

never positive

sometimes positive

Write the chemical equation representing Lattice enthalpy ($\Delta_{\text{lattice}}H$) of MgCl_2

Draw the energy diagram representing Lattice enthalpy ($\Delta_{\text{lattice}}H$)

- Using your own words define enthalpy of aquation ($\Delta_{\text{aq}}H$) :

- Is the enthalpy of aquation ($\Delta_{\text{aq}}H$) (choose one):

always positive

never positive

sometimes positive

Write the chemical equation representing enthalpy of aquation ($\Delta_{\text{aq}}H$) of MgCl_2 ,

Draw the energy diagram representing enthalpy of aquation ($\Delta_{\text{aq}}H$)

Draw the picture of hydrated MgCl_2 ions by water.

5. What do you need to know to find Enthalpy of solution (ΔH_{soln})

6. Using your own words define enthalpy of solution ($\Delta_{\text{sol}}H$):

7. Is the enthalpy of solution ($\Delta_{\text{sol}}H$) (choose one):

always positive

never positive

sometimes positive

Write the chemical equation representing enthalpy of solution ($\Delta_{\text{sol}}H$) of solid MgCl_2 ,

8. Can Enthalpy of solution (ΔH_{soln}) predict solubility?

1. Units of concentrations

a. Molality, m (mol/kg) = $\frac{\text{moles}(\text{solute})}{\text{mass}(\text{solvent})\text{kg}}$

2. Colligative property where i is a *van't Hoff factor*

a. Osmotic Pressure: $\Pi = MRT \cdot i$

b. Freezing Point Lowering : $\Delta T_{\text{fp}} = k_{\text{fp}} \cdot m \cdot i$

c. Boiling Point Elevation : $\Delta T_{\text{bp}} = k_{\text{bp}} \cdot m \cdot i$

d. Vapor Pressure Lowering: $p_{\text{solvent}} = \chi_{\text{solvent}} \cdot p^{\circ}_{\text{pure solv}}$; $\Delta p_{\text{solv}} = \chi_{\text{solute}} \cdot p^{\circ}_{\text{pure solv}}$

3. Refer to Page 466-478 in the text book Mahaffy

9. Is the freezing temperature of the solution always _____ compared to freezing temperature of pure solvent.

greater

lower

same


10. Is the boiling temperature of the solution always _____ compared to boiling temperature of pure solvent.

greater

lower

same

11. The figure below is the experimental plot of freezing point temperature, T_f , versus molality, m , for the solution of an unknown solute in cyclohexane. Calculate to two significant figures the freezing point depression constant K_f for cyclohexane, in $^{\circ}\text{C kg/mol}$. (use $i=1$ for this problem) (Answer: 20)



12. What is the boiling point of a 0.100*m* sucrose/water solution? ($K_{b,\text{water}} = 0.500 \text{ }^\circ\text{C}/m$) (Answer: 100.05)
13. Circle the solution below that has the lowest freezing point and explain your answer.
- 4mol NaCl(s) in 1L of water
 - 3mol CaCl₂(s) in 1L of water
 - 2mol AlCl₃(s) in 1L of water
 - 4mol C₆H₁₂O₆ (s) in 1L of water
14. The normal freezing point of cyclohexane is 6.6°C. A 0.2 g sample of an unknown solute is dissolved in 50 mL of cyclohexane ($d = 0.8 \text{ g/mL}$, $K_f = 20^\circ\text{C}/m$). If the freezing point of the solution is 3.6°C, what is the molar mass of the solute? (Answer: 30g/mol)
15. At 25 °C vapor pressure of pure water is 24 mmHg and that of seawater is 23.20 mmHg. Assuming that seawater contains only NaCl estimate its molal concentration.(Answer: 0.95)
16. The freezing point of benzene (C₆H₆) is 6 °C at 1 atm. In a laboratory experiment, students synthesized a new compound and found that when 40. grams of the compound were dissolved in 2000 grams of benzene ($K_f = 5.12 \text{ K}\cdot\text{kg}/\text{mol}$), the solution froze at 4 °C. The compound was also found to be nonvolatile and a non-electrolyte. Calculate the molar mass of the compound.(Answer:50)
17. 29.0 g of NaCl(s) is mixed with 200. g of ice. Calculate the boiling point of the mixture. The boiling point elevation constant for water is $K_b = 0.5^\circ\text{C}/\text{mol}$ (use $i=2$ for this problem) (Answer: 102.5)
18. What mass of CaCl₂ (MW = 111 g/mol; $i=3$) is required to lower the freezing point of 10.0 mL of water ($K_f=1.86^\circ\text{C}/m$) by 1°C?(Answer: 0.199)
19. Benzene freezes at 5.51°C ($K_f = 5.12 \text{ }^\circ\text{C}/m$). A benzene/acetic acid solution freezes at 2.01°C. How many moles of acetic acid were added to 500. g of benzene?(Answer:0.342)