

3. A 0.1M solution of NaX has $\text{pH} = 9.0$ at room temperature.
- Based on the information given do you have a reaction of acid with water or base with water? Why?
 - When salts that also contain an acid or base, like NaX, dissolve in water they ionize before they react. Write the reaction for NaX dissolving in water.
 - Write the chemical equation for the acid-base reaction that occurs in this solution. What are the conjugate acid and base pairs in the solution? NEVER include spectator ions in equilibrium reactions
 - Write the expression for equilibrium constant K (does your expression represent K_a or K_b):
 - What percent of the base reacted in solution? Is this a strong or weak base?
 - Based on the information given will you expect the equilibrium constant to be: (choose one)
 $K_b \ll 1$ or $K_b \gg 1$
4. Using the equilibrium constants in questions 2 and 3 derive the expression for equilibrium constant for autoionization of water.
- Write the chemical equation for autoionization of water:
 - Write the expression for equilibrium constant K_w :
 - Use questions 2c and 3d to derive the expression for K_w using K_a and K_b

5. A 0.1 M solution of an acid, HB, is found to have a pH of 3.52. (This suggests that it is an equilibrium pH). (Answer: 9×10^{-7} ; 0.3)

- a. Based on the information given do you have a reaction of acid with water or base with water? Why?

What is the chemical reaction?

Calculate equilibrium concentration of hydronium:

What is the equilibrium **expression** for K :

ICE table:

What was the percent reaction of the acid?

Calculate the value of K_a :

6. 0.5 M solution of NaD (D^- is a salt of the weak acid HD with the $K_a = 2.5 \times 10^{-10}$). (Answer: 11.6)

- a. Based on the information given do you have a reaction of acid with water or base with water? Why?

Write down net chemical reaction:

What is the equilibrium **expression**? (Is that a K_a , K_b or neither?)

Calculate the **value** of K for the reaction of D^- and water? (Hint: use the connection between K_a , K_b and K_w)

ICE table:

Fill in the data from the ICE table into your equilibrium **expression for K_b** :

pOH(at equilibrium) =

pH(at equilibrium) =

7. **Equal volumes** of a 0.10 M solution of a weak acid, HC, with $K_a = 1 \cdot 10^{-6}$, and a 0.20 M solution of NaOH are combined. What is the pH of the resulting solution? (Hint: what will react first? What will react completely? What is the limiting reagent?) (Answer: pH=12.7)
8. When 0.1mol of NaA is dissolved in 1L of pure water at room temperature, the pH is measured to be 7.0. Is HA a strong or weak acid?
- a. Based on the information given will you expect the equilibrium constant to be: (choose one)
- $K_a \ll 1$ or $K_a \gg 1$
9. Rank the acids HA, HB, HC, and HD from questions 5 through 8 in order of increasing acid strength.

10. You have a 1M solution of each of the salts below. Which of the salts will give the highest pH?
Which of the salts will give the lowest pH?

$\text{NH}_4\text{CH}_3\text{COO}$ (ammonium acetate),
 NH_4CN (ammonium cyanide),
 $\text{NH}_4\text{HC}_2\text{O}_4$ (ammonium oxalate).

	K_a	K_b
CH_3COOH	$2 \cdot 10^{-5}$	
HCN	$6 \cdot 10^{-10}$	
$\text{H}_2\text{C}_2\text{O}_4$	$6 \cdot 10^{-2}$	
NH_3		$2 \cdot 10^{-5}$

11. 250. mL of an 0.8 M solution of NaAc (salt of the weak acid HAc with the K_a of 2.5×10^{-10}) is added to 250. mL of water. (Answer: pH=11.6)

What is the Chemical Reaction?

What will be the concentration of Ac^- after dilution?

What is the Equilibrium Expression of K (Is it a K_a or K_b ?) for the reaction of NaD and water:

Find the value of the K:

Based on the value of K_b is D^- a strong base or a weak base?

Based on the value of K_a is HD a strong acid or a weak acid?

ICE table:

pOH(at equilibrium) =

pH(at equilibrium) =

Really Challenging problems to do at home:

12. One liter solution has 4.4 mol of HCl, 4.6 mol of NaOH and 0.2 mol of HCN. What is the pH when the resulting solution reaches equilibrium? (K_a of HCN is 5×10^{-10}) (Hint: what will react first? What will react completely? What is the limiting reagent?) (Answer: 11.3)

13. If equal volumes of 4.4 M HCl, 4.6 M NaOH and 0.2 M HCN are mixed (K_a of HCN is 5×10^{-10}), what is the pH when the resulting solution reaches equilibrium? (Answer: 11.1)

14. Determine the K_b of a base, at 25°C , if a 0.02 M aqueous solution of the base has a pH of 7.60 (This implies that it is an equilibrium pH). (Hint: compare pH of the solution with the pH of the water and decide if you can ignore the initial concentration of the H_3O^+ .) (Answer: 6×10^{-12})

Do at home to prepare for next week's discussion section.

1. You add HCl to a solution of equal moles of a weak acid and its conjugate base, and the number of moles of strong acid added is *smaller* than the number of moles of conjugate base initially present in the solution. Circle the correct answer(s) of the choices in the brackets.
 - a. The number of moles of A^- [**decreases / increases**] from the initial amount.
 - b. The number of moles of **HA** [**decreases / increases**] from the initial amount.
 - c. The number of moles of [**HCl / HA / A^-**] = 0 after neutralization reaction took place but before an equilibrium is established because [**HCl / HA / A^-**] is the limiting reagent and will be used up completely.
 - d. When the strong acid has completely reacted with the solution, you have a [**weak acid / weak base / strong acid / both weak acid and weak base**] present.

2. You add an HCl to a solution of equal moles of a weak acid and its conjugate base and the number of moles of strong acid added is *equal to* the number of moles of conjugate base initially present in the solution. Circle the correct answer(s) of the choices in the brackets:
 - a. The number of moles of A^- [decreases / increases] from the initial amount.
 - b. The number of moles of HA [decreases / increases] from the initial amount.
 - c. The number of moles of [HCl / HA / A^-] = 0 after neutralization reaction took place but before an equilibrium is established because [HCl / HA / A^-] is the limiting reagent and will be used up completely.
 - d. When the strong acid has completely reacted with the solution, you still have a [Weak acid / weak base / strong acid / both weak acid and weak base] present.

3. You add an HCl to a solution of equal moles of a weak acid and its conjugate base, and finally the number of moles of strong acid added is *greater than* the number of moles of conjugate base present in the solution initially. Circle the correct answer(s) of the choices in the brackets:
 - a. The number of moles of A^- [decreases / increases] from the initial amount.
 - b. The number of moles of HA [decreases / increases] from the initial amount.
 - c. The number of moles of [HCl / HA / A^-] = 0 after neutralization reaction took place but before an equilibrium is established because [HCl / HA / A^-] is the limiting reagent and will be used up completely.
 - d. When the HCl has completely reacted with the solution, you still have a
[Weak acid / weak base / strong acid / strong base] present.

4. You add NaOH to a solution of equal moles of a weak acid and its conjugate base, and the number of moles of strong base added is **smaller** than the number of moles of conjugate base initially present in the solution. Circle the correct answer(s) of the choices in the brackets.
- The number of moles of A^- [decreases / increases] from the initial amount.
 - The number of moles of HA [decreases / increases] from the initial amount.
 - The number of moles of $[OH^- / HA / A^-] = 0$ after neutralization reaction took place but before an equilibrium is established because $[OH^- / HA / A^-]$ is the limiting reagent and will be used up completely.
 - When NaOH has completely reacted with the solution, you still have a [Weak acid / weak base / strong acid / strong base/both weak acid and weak base] present.
5. You add a NaOH to a solution of equal moles of a weak acid and its conjugate base and the number of moles of strong base added is **equal to** the number of moles of conjugate base initially present in the solution. Circle the correct answer(s) of the choices in the brackets:
- The number of moles of A^- [decreases / increases] from the initial amount.
 - The number of moles of HA [decreases / increases] from the initial amount.
 - The number of moles of $[OH^- / HA / A^-] = 0$ after neutralization reaction took place but before an equilibrium is established because $[OH^- / HA / A^-]$ is the limiting reagent and will be used up completely.
 - When NaOH has completely reacted with the solution, you still have a [weak acid / weak base / strong acid / strong base/both weak acid and weak base] present.
6. You add a NaOH to a solution of equal moles of a weak acid and its conjugate base, and finally the number of moles of strong base added is **greater than** the number of moles of conjugate base present in the solution initially. Circle the correct answer(s) of the choices in the brackets:
- The number of moles of A^- [decreases / increases] from the initial amount.
 - The number of moles of HA [decreases / increases] from the initial amount.
 - The number of moles of $[OH^- / HA / A^-] = 0$ after neutralization reaction took place but before an equilibrium is established because $[OH^- / HA / A^-]$ is the limiting reagent and will be used up completely.
 - When the NaOH has completely reacted with the solution, you still have a [weak acid / weak base / strong acid / strong base] present.